



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ferred to a dish of cold water or alcohol standing at the end of the embedding stage and into which it is immersed as soon as the paraffine is cooled sufficiently to prevent the breaking of the surface film by the water.

The use of this embedding stage makes unnecessary the use of the top of the paraffine bath for this purpose. Its use helps greatly in securing good embedding because it permits the paraffine to be melted clear to the bottom of the embedding tray and thus the orientation is made easy. The plan of allowing the object to lie on a layer of congealed paraffine is not only unnecessary but is faulty in that the paraffine is too soft to permit accurate orientation of the object and also because a cleavage plane is formed at which the paraffine frequently breaks during the sectioning. The stage is convenient to use, does away with the necessity of using gas, and largely obviates the danger of overheating the tissue which danger is always present when a gas flame is used for heating the ordinary stage. This stage because of its low construction is very stable, unlike the very insecure stage used with the gas flame, and with the levelling screws it may be levelled. In several months' use by a class no objectionable features have appeared and its good points are only the better appreciated.

*Zoölogical Laboratory,
University of Michigan.*

GEORGE R. LA RUE.

MAKING GLASS PLATES FOR COVERING MUSEUM JARS

At this time when it is impossible to secure from abroad the glass plates for covering museum jars it is worth while to know that after a little practise passable plates may be made in any laboratory equipped with power grinding and buffing machinery. Double strength glass plates may be purchased cut to size or they may be cut in the laboratory. Their edges may be rounded and a narrow ground surface at the margin may be secured by grinding on a carborundum wheel designated 120J-G5 which can be purchased from the Carborundum Co., Niagara Falls. The size of the wheel will depend somewhat on the power and speed of the grinder. In this laboratory a $4\frac{1}{2}$ by $\frac{1}{2}$ inch wheel belted to a $\frac{1}{2}$ h. p. motor

of about 1800 R. P. M. is used. While plates made in this way are not as good as the imported article they are usable and cheap, and by this means museum jars whose covers have been broken may be put into use.

*Zoölogical Laboratory,
University of Michigan.*

GEORGE R. LA RUE.

THE POSSIBLE NATURE OF THE "BOOK LUNGS" OF SPIDERS

The abdomen of spiders is now unsegmented, and yet it is probable that spiders have descended from ancestors whose bodies were segmented throughout.

The breathing apparatus in spiders is varied, some forms showing some development of tracheal tubes. On the forward end of the abdomen are found two sacs, each of which encloses a folded membrane which exposes the blood to the air. These are the book lungs.

In the section of such a lung from an *Aglena* (Plate XX, Fig. 1) the membranous character of the organ will be seen. Red blood cells may be seen between the double membranes. The outer surface of the membrane is covered with short spines, which prevent the moist membranes from adhering.

It is possible that this arrangement is derived from an ancestral form which had external gills at this point, somewhat similar to the tracheal gill membranes of insect nymphs.

A figure of a section of the young wing membranes of an *Ephemera* nymph is shown (Pl. XX, Fig. 2) for comparison. The similarity of structure is striking.

E. W. ROBERTS.

NOTE ON THE NATURE OF THE CYTO-PLASTID

The cyto-plasm of a cell contains unit plastids which themselves bear a great resemblance to a complete cell with its nucleus and cyto-plasm.

Using the Tussock Moth egg for an illustration we get a suggestion of this condition. The egg is filled with nutritive material supplied by numerous nurse cells from their own cyto-some system.